



COOPER SPECIALTY FEEDS



Show Goat Barley Loco 17%

A complete texturized feed for growth and finish of goats for show competition

| Key Features | Benefits |
|--|---|
| Steam flaked corn & barley | <ul style="list-style-type: none">Improves digestibility and reduces fines |
| Roasted soybeans | <ul style="list-style-type: none">Excellent source of bypass protein and fat |
| Multiple degradable and bypass protein sources | <ul style="list-style-type: none">Helps support rumen fermentation, muscle growth and expression |
| Yeast Culture | <ul style="list-style-type: none">Supports the immune system and rumen fermentation |
| Ammonium Chloride | <ul style="list-style-type: none">To help prevent urinary calculi |
| Trace Mineral Amino Acid Complexes | <ul style="list-style-type: none">Increases bioavailability for improved overall health, hoof and hair coat quality |
| Rumensin® (monensin) | <ul style="list-style-type: none">For the prevention of coccidiosis |

| Guaranteed Analysis | |
|------------------------|--------------|
| Active Drug Ingredient | |
| Monensin | 20 grams/ton |
| Crude Protein, min | 17.0% |
| NPN, max | 1.5% |
| Crude Fat, min | 5.0% |
| Crude Fiber, max | 10.0% |
| Calcium, min | 0.7% |
| Calcium, max | 1.2% |
| Phosphorus, min | 0.40% |
| Copper, min | 5.0 ppm |
| Copper, max | 10.0 ppm |
| Selenium, min | 0.30 ppm |
| Vitamin A, min | 4000 IU/lb |

Feeding Instructions

- Feed continuously as sole ration to goats. Provide a constant supply of clean water along with a small quantity of long-stemmed hay.
- CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin medicated goat feeds are safe for use in goats only. Consumption by unapproved species may result in toxic reactions. Do not feed to lactating goats.

www.cooperfeeds.com

Feed2Win@cooperfeeds.com